

Whitney type inequalities for local anisotropic polynomial approximation

Dinh Dung^{a,*}; Tino Ullrich^b

^a Vietnam National University, Hanoi, Information Technology Institute
144, Xuan Thuy, Hanoi, Vietnam

^bHausdorff-Center for Mathematics, 53115 Bonn, Germany

June 6, 2011

Abstract

We prove a multivariate Whitney type theorem for the local anisotropic polynomial approximation in $L_p(Q)$ with $1 \leq p \leq \infty$. Here Q is a d -parallelepiped in \mathbb{R}^d with sides parallel to the coordinate axes. We consider the error of best approximation of a function f by algebraic polynomials of fixed degree at most $r_i - 1$ in variable x_i , $i = 1, \dots, d$, and relate it to a so-called total mixed modulus of smoothness appropriate to characterizing the convergence rate of the approximation error. This theorem is derived from a Johnen type theorem on equivalence between a certain K-functional and the total mixed modulus of smoothness which is proved in the present paper.

Keywords Whitney type inequality; Anisotropic approximation by polynomials; Total mixed modulus of smoothness; Mixed K-functional; Sobolev space of mixed smoothness;

Mathematics Subject Classifications (2000) 41A10; 41A50; 41A63

1 Introduction and main results

The classical Whitney theorem establishes the equivalence between the modulus of smoothness $\omega_r(f, |I|)_{p,I}$ and the error of best approximation $E_r(f)_{p,I}$ of a function $f : I \rightarrow \mathbb{R}$ by algebraic polynomials of degree at most $r - 1$ measured in L_p , $1 \leq p \leq \infty$, where $I := [a, b]$ is an interval in \mathbb{R} and $|I| = b - a$ its length. Namely, the following inequalities

$$2^{-r} \omega_r(f, |I|)_{p,I} \leq E_r(f)_{p,Q} \leq C \omega_r(f, |I|)_{p,I} \quad (1.1)$$

hold true with a constant C depending only on r . This result was first proved by Whitney [24] for $p = \infty$ and extended by Brudnyĭ [2] to $1 \leq p < \infty$. The inequalities (1.1) provide, in particular,

*Corresponding author. Email: dinhzung@gmail.com

a convergence characterization for a local polynomial approximation when the degree $r - 1$ of polynomials is fixed and the interval I is small.

Several authors have dealt with this topic in order to extend and generalize the result in various directions. Let us briefly mention them. A multivariate (isotropic) generalization for functions on a coordinate d -cube Q in \mathbb{R}^d was given by Brudnyĭ [3] and [4]. It turned out that the result keeps valid if one replaces the d -cube by a more general domain Ω . The case of a convex domain $\Omega \subset \mathbb{R}^d$ is already treated in Brudnyĭ [3]. Let us also refer to the recent contributions by Dekel, Leviatan [7] and Dekel [6] with focus on convex and Lipschitz domains and the improvement of the constants involved.

A reasonable question is also to ask for the case $0 < p < 1$. We refer to the works of Storozhenko [18], Storozhenko, Oswald [19], and in addition to the appendix of the substantial paper by Hedberg and Netrusov [12] for a brief history and further references.

A natural question arises: Is there a Whitney type theorem for the anisotropic approximation of multivariate functions on a coordinate d -parallelepiped Q ? Some work has been done in this direction, see for instance Garrigós, Tabacco [13]. However, the present paper deals with a rather different setting which is somehow related to the theory of function spaces with mixed smoothness properties [10, 20, 22, 23]. We intend to approximate a multivariate function f by polynomials of fixed degree at most $r_i - 1$ in variable x_i , $i = 1, \dots, d$, on a small d -parallelepiped Q . A total mixed modulus of smoothness is defined which turns out to be a suitable convergence characterization to this approximation. The classical Whitney inequality can be derived as a corollary of Johnen's theorem [14] on the equivalence of the r -th Peetre K -functional $K_r(f, t^r)_{p, I}$, see [16], and the modulus of smoothness $\omega_r(f, t)_{p, I}$. A proof was given by Johnen and Scherer in [15]. Following this approach to Whitney type theorems we will introduce the notion of a mixed K -functional and prove its equivalence to the total mixed modulus of smoothness by generalizing the technique of Johnen and Scherer to the multivariate mixed situation.

1.1 Notation

In order to give an exact setting of the problem and formulate the main results let us preliminarily introduce some necessary notations. As usual, \mathbb{N} is reserved for the natural numbers, by \mathbb{Z} we denote the set of all integers, and by \mathbb{R} the real numbers. Furthermore, \mathbb{Z}_+ and \mathbb{R}_+ denote the set of non-negative integers and real numbers, respectively. Elements x of \mathbb{R}^d will be denoted by $x = (x_1, \dots, x_d)$. For a vector $r \in \mathbb{Z}_+^d$ and $x \in \mathbb{R}^d$, we will further write

$$x^r := (x_1^{r_1}, \dots, x_d^{r_d}).$$

Moreover, if $x, y \in \mathbb{R}^d$, the inequality $x \leq y$ ($x < y$) means that $x_i \leq y_i$ ($x_i < y_i$), $i = 1, \dots, d$. As usual, the notation $A \ll B$ indicates that there is a constant $c > 0$ (independent of the parameters which are relevant in the context) such that $A \leq cB$, whereas $A \asymp B$ is used if $A \ll B$ and $B \ll A$, respectively.

If $r \in \mathbb{N}^d$, let \mathcal{P}_r be the set of algebraic polynomials of degree at most $r_i - 1$ at variable x_i , $i \in [d]$, where $[d]$ denotes the set of all natural numbers from 1 to d . We intend to approximate a function f defined on a d -parallelepiped

$$Q := [a_1, b_1] \times \dots \times [a_d, b_d]$$

by polynomials from the class \mathcal{P}_r . If $D \subset \mathbb{R}^d$ is a domain in \mathbb{R}^d we denote by $L_p(D)$, $0 < p \leq \infty$, the quasi-normed space of Lebesgue measurable functions on D with the usual p -th integral quasi-norm $\|\cdot\|_{p,D}$ to be finite, whereas we use the ess sup norm if $p = \infty$. The error of the best approximation of $f \in L_p(Q)$ by polynomials from \mathcal{P}_r is measured by

$$E_r(f)_{p,Q} := \inf_{\varphi \in \mathcal{P}_r} \|f - \varphi\|_{p,Q}.$$

For $r \in \mathbb{Z}_+$, $h \in \mathbb{R}$, and a univariate functions f , the r th difference operator Δ_h^r is defined by

$$\Delta_h^r(f, x) := \sum_{j=0}^r (-1)^{r-j} \binom{r}{j} f(x + jh) \quad , \quad \Delta_h^0 f(x) := f(x),$$

whereas for $r \in \mathbb{Z}_+^d$, $h \in \mathbb{R}^d$ and a d -variate function $f : \mathbb{R}^d \rightarrow \mathbb{R}$, the mixed r th difference operator Δ_h^r is defined by

$$\Delta_h^r := \prod_{i=1}^d \Delta_{h_i, i}^{r_i}.$$

Here, the univariate operator $\Delta_{h_i, i}^{r_i}$ is applied to the univariate function f by considering f as a function of variable x_i with the other variables fixed. Let

$$\omega_r(f, t)_{p,Q} := \sup_{|h_i| \leq t_i, i \in [d]} \|\Delta_h^r(f)\|_{p, Q_{rh}} \quad , \quad t \in \mathbb{R}_+^d,$$

be the mixed r th modulus of smoothness of f , where for $y, h \in \mathbb{R}^d$, we write $yh := (y_1 h_1, \dots, y_d h_d)$ and $Q_y := \{x \in Q : x_i, x_i + y_i \in [a_i, b_i], i \in [d]\}$. For $r \in \mathbb{Z}_+^d$ and $e \subset [d]$, denote by $r(e) \in \mathbb{Z}_+^d$ the vector with $r(e)_i = r_i, i \in e$ and $r(e)_i = 0, i \notin e$ ($r(\emptyset) = 0$). If $r \in \mathbb{N}^d$, we define the *total mixed modulus of smoothness of order r* by

$$\Omega_r(f, t)_{p,Q} := \sum_{e \subset [d], e \neq \emptyset} \omega_{r(e)}(f, t)_{p,Q}, \quad t \in \mathbb{R}_+^d.$$

This particular modulus of smoothness is not new. In the periodic context the total mixed modulus of smoothness $\Omega_r(f, \cdot)_{\infty, Q}$ has been used in [5] for estimations of the convergence rate of the approximation of continuous periodic functions by rectangular Fourier sums. Moreover, $\Omega_r(f, \cdot)_{p, Q}$ is related to mixed moduli of smoothness necessary for characterizing function spaces with dominating mixed smoothness properties, see [10, 20] and the recent contributions [22, 23, 21, 11].

1.2 Main results

In the present paper, we generalize the Whitney inequality (1.1) to the error of the best local anisotropic approximation $E_r(f)_{p,Q}$ by polynomials from \mathcal{P}_r and the total mixed modulus of smoothness $\Omega_r(f, t)_{p,Q}$. More precisely, we prove the following Whitney type inequalities.

Theorem 1.1 *Let $1 \leq p \leq \infty$, $r \in \mathbb{N}^d$. Then there is a constant C depending only on r, d such that for every $f \in L_p(Q)$*

$$\left(\sum_{e \subset [d]} \prod_{i \in e} 2^{r_i} \right)^{-1} \Omega_r(f, \delta)_{p, Q} \leq E_r(f)_{p, Q} \leq C \Omega_r(f, \delta)_{p, Q}, \quad (1.2)$$

where $\delta = \delta(Q) := (b_1 - a_1, \dots, b_d - a_d)$ is the size of Q .

Theorem 1.1 shows that the total mixed modulus of smoothness $\Omega_r(f, t)_{p, Q}$ gives a sharp convergence characterization of the best anisotropic polynomial approximation when r is fixed and the size $\delta(Q)$ of the d -parallelepiped Q is small. This may have applications in the approximation of functions with mixed smoothness by piecewise polynomials or splines.

So far we focus on the case $1 \leq p \leq \infty$. This makes it possible to apply a technique developed by Johnen and Scherer [15]. As mentioned above they showed the equivalence of Peetre's K -functional of order r with respect to a classical Sobolev space W_p^r and the modulus of smoothness of order r for the univariate case. The question of a K -functional suitable for mixed Sobolev spaces has been often considered in the past. We refer for instance to Sparr [17] and DeVore et al. [9]. By introducing a mixed K -functional $K_r(f, t)_{p, Q}$, $t \in \mathbb{R}_+^d$, (see the definition in Section 3), such an equivalence between $K_r(f, t^r)_{p, Q}$ and the total mixed modulus of smoothness $\Omega_r(f, t)_{p, Q}$ can be established as well. Namely, we prove the following

Theorem 1.2 *Let $1 \leq p \leq \infty$ and $r \in \mathbb{N}^d$. Then for any $f \in L_p(Q)$, the following inequalities*

$$\left(\sum_{e \subset [d]} \prod_{i \in e} 2^{r_i} \right)^{-1} \Omega_r(f, t)_{p, Q} \leq K_r(f, t^r)_{p, Q} \leq C \Omega_r(f, t)_{p, Q} \quad , \quad t \in \mathbb{R}_+^d, \quad (1.3)$$

hold true with a constants C depending on r, p, d only.

The paper is organized as follows. In Section 2 we establish an error estimate for the anisotropic polynomial approximation for functions from Sobolev spaces of mixed smoothness. Section 3 is devoted to the equivalence of the total mixed modulus of smoothness and the mixed K -functional (Theorem 1.2) which is applied in Section 4 to derive the Whitney type inequality for the local anisotropic polynomial approximation (Theorem 1.1).

2 Anisotropic polynomial approximation in Sobolev spaces of mixed smoothness

By $f^{(k)}$, $k \in \mathbb{Z}_+^d$, we denote the k th order generalized mixed derivative of a locally integrable function f , i.e.,

$$\int_Q f^{(k)}(x) \varphi(x) dx = (-1)^{k_1 + \dots + k_d} \int_Q f(x) \frac{\partial^{k_1 + \dots + k_d} \varphi}{\partial x_1^{k_1} \dots \partial x_d^{k_d}}(x) dx$$

for all test functions $\varphi \in C_0^\infty(Q)$, where $C_0^\infty(Q)$ is the space of infinitely differentiable functions on Q with compact support, which is interior to Q . If a function f possesses sth locally integrable classical partial derivatives for all $s \leq k$ on Q , then the k th generalized derivative of f coincides with the k th classical partial derivative. In this case, we identify both and use the same notation $f^{(k)}$.

For $r \in \mathbb{Z}_+^d$ and $1 \leq p \leq \infty$, the Sobolev space $W_p^r(Q)$ of mixed smoothness r is defined as the set of functions $f \in L_p(Q)$, for which the generalized derivative $f^{(r(e))}$ exists as a locally integrable function for all $e \subset [d]$, and the following norm is finite

$$\|f\|_{W_p^r(Q)} := \sum_{e \subset [d]} \|f^{(r(e))}\|_{p,Q}.$$

We aim at giving an upper bound of the error of best approximation of $f \in W_p^r(Q)$ by polynomials of degree $r_i - 1$ with respect to the variable x_i , $i = 1, \dots, d$. For this purpose we need some auxiliary lemmas. To begin with we deal with univariate functions. The following lemma is proven in [8, page 38].

Lemma 2.1 *Let $1 \leq p \leq \infty$, $r \geq 1$ and $Q = [a, b]$. Then there exist constants C_1, C_2 depending only on r such that for $k = 0, \dots, r - 1$ and $0 \leq t \leq b - a$ the inequality*

$$t^k \|f^{(k)}\|_{p,Q} \leq C_1 (\|f\|_{p,Q} + t^r \|f^{(r)}\|_{p,Q}), \quad (2.1)$$

holds true for any $f \in W_p^r(Q)$.

Lemma 2.2 *Let $r \in \mathbb{Z}_+^d$, $1 \leq p \leq \infty$, and $Q = [0, b_1] \times \dots \times [0, b_d]$ where $b_i > 0$, $i = 1, \dots, d$. For fixed $f \in W_p^r(Q)$, $k \leq r$, and $j \in [d]$ the univariate function*

$$g := f^{(k - k_j e_j)}(x_1, \dots, x_{j-1}, \cdot, x_{j+1}, \dots, x_d)$$

belongs to $W_p^{r_j}([0, b_j])$ for almost all $x_i \in [0, b_i]$, $i \in [d] \setminus \{j\}$.

Proof. Let $\varphi_i \in C_0^\infty(0, b_i)$, $i = 1, \dots, d$, be arbitrary smooth compactly supported functions. Clearly, the tensor product $\Phi(x_1, \dots, x_d) := \prod_{i \in [d]} \varphi_i(x_i)$ belongs to $C_0^\infty(Q)$. Then, for $0 \leq \ell_j \leq r_j$

$$\begin{aligned} & \int_0^{b_1} \dots \int_0^{b_{j-1}} \int_0^{b_{j+1}} \dots \int_0^{b_d} \prod_{\substack{i \in [d] \\ i \neq j}} \varphi_i(x_i) \left(\int_0^{b_j} f^{(k - k_j e_j)}(x_1, \dots, x_{j-1}, t, x_{j+1}, \dots, x_d) \varphi_j^{(\ell_j)}(t) dt \right) \prod_{\substack{i \in [d] \\ i \neq j}} dx_i \\ &= \int_0^{b_1} \dots \int_0^{b_d} f^{(k - k_j e_j)}(x_1, \dots, x_d) \Phi^{(0, \dots, \ell_j, 0, \dots, 0)}(x_1, \dots, x_d) dx_1 \dots dx_d \\ &= (-1)^{\ell_j} \int_0^{b_1} \dots \int_0^{b_d} f^{(k + e_j(\ell_j - k_j))}(x_1, \dots, x_d) \Phi(x_1, \dots, x_d) dx_1 \dots dx_d \\ &= \int_0^{b_1} \dots \int_0^{b_{j-1}} \int_0^{b_{j+1}} \dots \int_0^{b_d} \prod_{\substack{i \in [d] \\ i \neq j}} \varphi_i(x_i) \left(\int_0^{b_j} f^{(k + e_j(\ell_j - k_j))}(x_1, \dots, x_{j-1}, t, x_{j+1}, \dots, x_d) \varphi_j(t) dt \right) \prod_{\substack{i \in [d] \\ i \neq j}} dx_i. \end{aligned}$$

This implies the coincidence of the dt -integrals in the first and last line almost everywhere (with respect to x_i , $i \in [d] \setminus \{j\}$). Therefore, the generalized derivatives of order ℓ_j exist as a locally integrable function, in fact, they coincide with $f^{(k+e_j(\ell_j-k_j))}(x_1, \dots, x_{j-1}, \cdot, x_{j+1}, \dots, x_d)$. This is a function from $L_p([0, b_j])$ (almost everywhere with respect to x_i) since f belongs to $W_p^r(Q)$. Therefore, we have $g \in W_p^{r_j}([0, b_j])$. \square

The following result is interesting on its own. It generalizes the content of [8, Theorem 5.3] to the multivariate situation. The statement is not very surprising and probably known. However, since we did not find a proper reference in the literature, a proof is provided.

Lemma 2.3 *Let $r \in \mathbb{N}^d$ and $Q = [0, b_1] \times \dots \times [0, b_d]$. Let further $f \in L_1(Q)$ such that $f^{(r(e))} = 0$ for all non-empty subsets $e \subset [d]$. Then f coincides almost everywhere with a polynomial P of degree $r - 1$, i.e., $f \in \mathcal{P}_r$.*

Proof. For simplicity reasons we give a proof for $d = 2$, so let $Q = [0, b_1] \times [0, b_2]$. We follow the inductive argument in the proof of the corresponding one-dimensional statement [8, Theorem 5.3]. The latter and Lemma 2.2 imply the statement in case $r = (1, 1)$. Assume now that it is proven for some $r \in \mathbb{N}^2$. Put $\bar{r} = (r_1 + 1, r_2)$ without loss of generality. We will prove that the assumption

$$f^{(\bar{r}(e))} = 0 \text{ for all non-empty subsets } e \subset [d] \quad (2.2)$$

implies that f coincides almost everywhere with a polynomial $P \in \mathcal{P}_{\bar{r}}$. To do this we need to construct special test functions. Choose a function $\psi \in C_0^\infty(Q)$ arbitrarily and let $h \in C_0^\infty([0, b_1])$ be a univariate function such that $\int_0^{b_1} h(t)dt = 1$. We define the functions

$$\begin{aligned} \varphi(x_1, x_2) &:= \psi(x_1, x_2) - h(x_1) \int_0^{b_1} \psi(s, x_2) ds, \\ \Phi(x_1, x_2) &:= \int_0^{x_1} \varphi(s, x_2) ds. \end{aligned} \quad (2.3)$$

This construction gives immediately $\Phi \in C_0^\infty(Q)$. By our assumption (2.2) we have in particular

$$\begin{aligned} 0 &= \int_Q \Phi^{(r_1+1, 0)} f dx_1 dx_2 \\ &= \int_Q \psi^{(r_1, 0)} f dx_1 dx_2 - \int_Q f(x_1, x_2) h^{(r_1)}(x_1) \int_{s=0}^{b_1} \psi(s, x_2) ds dx_1 dx_2 \\ &= \int_Q \psi^{(r_1, 0)} f dx_1 dx_2 - \int_Q f(x_1, x_2) h^{(r_1)}(x_1) \int_{s=0}^{b_1} \psi^{(r_1, 0)}(s, x_2) \frac{s^{r_1}}{r_1!} ds dx_1 dx_2 \\ &= \int_{s=0}^{b_1} \int_{x_2=0}^{b_2} \psi^{(r_1, 0)}(s, x_2) \cdot \left(f(s, x_2) - \frac{s^{r_1}}{r_1!} \int_{x_1=0}^{b_1} f(x_1, x_2) h^{(r_1)}(x_1) dx_1 \right) dx_2 ds \\ &= \int_Q \psi^{(r_1, 0)}(s, x_2) \cdot \left(f(s, x_2) - \frac{s^{r_1}}{r_1!} \int_{x_1=0}^{b_1} f(x_1, x_2) h^{(r_1)}(x_1) dx_1 \right) ds dx_2. \end{aligned} \quad (2.4)$$

Analogously we see

$$\begin{aligned}
0 &= \int_Q \Phi^{(r_1+1, r_2)} f dx_1 dx_2 \\
&= \int_Q \psi^{(r_1, r_2)}(s, x_2) \cdot \left(f(s, x_2) - \frac{s^{r_1}}{r_1!} \int_{x_1=0}^{b_1} f(x_1, x_2) h^{(r_1)}(x_1) dx_1 \right) ds dx_2.
\end{aligned} \tag{2.5}$$

Using (2.2) once more we get for any $s \in [0, b_1]$

$$\int_0^{b_1} \int_0^{b_2} h^{r_1}(x_1) \psi^{(0, r_2)}(s, x_2) f(x_1, x_2) dx_2 dx_1 = 0$$

which implies

$$0 = \int_Q \psi^{(0, r_2)}(s, x_2) \cdot \left(f(s, x_2) - \frac{s^{r_1}}{r_1!} \int_0^{b_1} f(x_1, x_2) h^{(r_1)}(x_1) dx_1 \right) ds dx_2. \tag{2.6}$$

Since ψ was chosen arbitrarily our induction hypothesis together with (2.4), (2.5), (2.6) imply that the function

$$g(s, x_2) := f(s, x_2) - \frac{s^{r_1}}{r_1!} \int_0^{b_1} f(x_1, x_2) h^{(r_1)}(x_1) dx_1$$

is a bivariate polynomial from \mathcal{P}_r . If we show that the univariate function

$$p(t) = \int_0^{b_1} f(x_1, t) h^{(r_1)}(x_1) dx_1$$

is a polynomial of degree at most $r_2 - 1$, we prove that $f \in \mathcal{P}_{\bar{r}}$. Indeed, let $\varphi \in C_0^\infty([0, b_2])$ arbitrary then

$$\int_0^{b_2} \varphi^{(r_2)}(t) p(t) dt = \int_Q f(x_1, x_2) h^{(r_1)}(x_1) \varphi^{(r_2)}(x_2) dx_1 dx_2 = 0 \tag{2.7}$$

by using (2.2) once more. This together with [8, Theorem 5.3] implies that p is a univariate polynomial of degree at most $r_2 - 1$. The proof is finished in case $d = 2$. For $d > 2$ the argument is essentially the same. Note that in this situation one needs an additional inductive step with respect to d to adapt the argument after (2.6). \square

By using the previous result we are now able to define a Taylor type polynomial via its integral representation. For simplicity we restrict again to the case $d = 2$. A corresponding statement holds true in case $d > 2$, too. See Remark 2.6 below.

Lemma 2.4 *Let $r \in \mathbb{N}^2$, $1 \leq p \leq \infty$, and $f \in W_p^r(Q)$ for $Q = [0, b_1] \times [0, b_2]$. Then the function $P_r f$ defined by*

$$\begin{aligned}
P_r f(x_1, x_2) &:= f(x_1, x_2) - \int_0^{x_2} f^{(0, r_2)}(x_1, t) \frac{(x_2 - t)^{r_2-1}}{(r_2 - 1)!} dt \\
&\quad - \int_0^{x_1} f^{(r_1, 0)}(s, x_2) \frac{(x_1 - s)^{r_1-1}}{(r_1 - 1)!} ds \\
&\quad + \int_0^{x_1} \int_0^{x_2} f^{(r_1, r_2)}(s, t) \frac{(x_1 - s)^{r_1-1}}{(r_1 - 1)!} \frac{(x_2 - t)^{r_2-1}}{(r_2 - 1)!} dt ds
\end{aligned} \tag{2.8}$$

is well-defined and coincides almost everywhere with a polynomial from \mathcal{P}_r .

Proof. Since f is from $W_p^r(Q)$, i.e., all the derivatives belong to $L_p(Q) \subset L_1(Q)$, the function $P_r f$ is well-defined. We intend to apply Lemma 2.3 in order to obtain $P_r f \in \mathcal{P}_r$. Let us compute the derivatives $(P_r f)^{(r_1,0)}$, $(P_r f)^{(0,r_2)}$, and $(P_r f)^{(r_1,r_2)}$. Choose $\varphi \in C_0^\infty(Q)$ arbitrarily. We start with $(P_r f)^{(r_1,0)}$. By changing the order of integration we get

$$\begin{aligned}
& \int_Q P_r f(x_1, x_2) \varphi^{(r_1,0)} dx_1 dx_2 = \int_Q f(x_1, x_2) \varphi^{(r_1,0)}(x_1, x_2) dx_1 dx_2 \\
& - \int_{t=0}^{b_2} \int_{x_1=0}^{b_1} f^{(0,r_2)}(x_1, t) \int_{x_2=t}^{b_2} \frac{(x_2-t)^{r_2-1}}{(r_2-1)!} \varphi^{(r_1,0)}(x_1, x_2) dx_2 dx_1 dt \\
& - \int_{x_2=0}^{b_2} \int_{s=0}^{b_1} f^{(r_1,0)}(s, x_2) \int_{x_1=s}^{b_1} \frac{(x_1-s)^{r_1-1}}{(r_1-1)!} \varphi^{(r_1,0)}(x_1, x_2) dx_1 ds dx_2 \\
& + \int_{s=0}^{b_1} \int_{t=0}^{b_2} f^{(r_1,r_2)}(s, t) \int_{x_1=s}^{b_1} \int_{x_2=t}^{b_2} \frac{(x_1-s)^{r_1-1}}{(r_1-1)!} \frac{(x_2-t)^{r_2-1}}{(r_2-1)!} \varphi^{(r_1,0)}(x_1, x_2) dx_2 dx_1 dt ds.
\end{aligned} \tag{2.9}$$

Integration by parts shows that

$$\int_{x_1=s}^{b_1} \frac{(x_1-s)^{r_1-1}}{(r_1-1)!} \varphi^{(r_1,0)}(x_1, x_2) dx_1 = (-1)^{r_1} \varphi(s, x_2) \tag{2.10}$$

and the third summand on the right-hand side of (2.9) can therefore be rewritten to

$$\begin{aligned}
& - \int_{x_2=0}^{b_2} \int_{s=0}^{b_1} f^{(r_1,0)}(s, x_2) \int_{x_1=s}^{b_1} \frac{(x_1-s)^{r_1-1}}{(r_1-1)!} \varphi^{(r_1,0)}(x_1, x_2) dx_1 ds dx_2 \\
& = -(-1)^{r_1} \int_{x_2=0}^{b_2} \int_{s=0}^{b_1} f^{(r_1,0)}(s, x_2) \varphi(s, x_2) ds dx_2 \\
& = - \int_{x_2=0}^{b_2} \int_{s=0}^{b_1} f(s, x_2) \varphi^{(r_1,0)}(s, x_2) ds dx_2
\end{aligned} \tag{2.11}$$

which cancels the first summand. Using (2.10) ones more we can rewrite the last summand in (2.9) to

$$\begin{aligned}
& (-1)^{r_1} \int_{s=0}^{b_1} \int_{t=0}^{b_2} f^{(r_1,r_2)}(s, t) \int_{x_2=t}^{b_2} \frac{(x_2-t)^{r_2-1}}{(r_2-1)!} \varphi(s, x_2) dx_2 dt ds \\
& = \int_{s=0}^{b_1} \int_{t=0}^{b_2} f^{(0,r_2)}(s, t) \int_{x_2=t}^{b_2} \frac{(x_2-t)^{r_2-1}}{(r_2-1)!} \varphi^{(r_1,0)}(s, x_2) dx_2 dt ds
\end{aligned} \tag{2.12}$$

which cancels the second summand. Hence, we obtain $(P_r f)^{(r_1,0)} = 0$ since φ was chosen arbitrarily. A similar effect occurs if we deal with $\int_Q P_r f(x_1, x_2) \varphi^{(0,r_2)} dx_1 dx_2$ which gives that also $(P_r f)^{(0,r_2)} = 0$. In case of $\int_Q P_r f(x_1, x_2) \varphi^{(r_1,r_2)} dx_1 dx_2$ we easily see that both the (modified) second and third summand in (2.9) can be rewritten to the negative of the first summand. However, the (modified) last summand can be rewritten to the first summand itself. Finally, all four summands sum up to zero. \square

Remark 2.5 The polynomial $P_r f$ in (2.8) can be identified with the bivariate Taylor polynomial

$$T_r f(x_1, x_2) := \sum_{k_2=0}^{r_2-1} \sum_{k_1=0}^{r_1-1} f^{(k_1, k_2)}(0, 0) \frac{x_1^{k_1}}{k_1!} \frac{x_2^{k_2}}{k_2!} \quad (2.13)$$

in the following sense. If $r \in \mathbb{N}^2$, $1 \leq p \leq \infty$, $Q = [0, b_1] \times [0, b_2]$, and $f \in W_p^r(Q)$ then f has continuous derivatives of order $k < r$. This result is implicitly contained in the book [1]. Indeed, it is a combination of multiparameter Sobolev averaging using product kernels in Section 2.7.10 and 3.13 with the estimates in 3.10, especially Theorem 3.10.4. The condition involving r and k there has to be replaced by the componentwise condition $k < r$. We omit details. Consequently, it makes sense to define the Taylor polynomial (2.13). Integration by parts shows that $T_r f$ coincides almost everywhere with $P_r f$ in (2.8). Hence, for functions from $W_p^r(Q)$ we have the Taylor formula

$$\begin{aligned} T_r f(x_1, x_2) &= f(x_1, x_2) - \int_0^{x_2} f^{(0, r_2)}(x_1, t) \frac{(x_2 - t)^{r_2-1}}{(r_2 - 1)!} dt \\ &\quad - \int_0^{x_1} f^{(r_1, 0)}(s, x_2) \frac{(x_1 - s)^{r_1-1}}{(r_1 - 1)!} ds \\ &\quad + \int_0^{x_1} \int_0^{x_2} f^{(r_1, r_2)}(s, t) \frac{(x_1 - s)^{r_1-1}}{(r_1 - 1)!} \frac{(x_2 - t)^{r_2-1}}{(r_2 - 1)!} dt ds. \end{aligned} \quad (2.14)$$

Remark 2.6 Lemma 2.4 and the Taylor formula (2.14) have an obvious counterpart in d dimensions. Note that the sum in (2.14) is twice the iteration (componentwise) of the one dimensional integral

$$T_r f(x) := f(x) - \int_0^x f^{(r)}(s) \frac{(x - s)^{r-1}}{(r - 1)!} ds. \quad (2.15)$$

The d -times iteration of this procedure results in a sum of iterated integrals where the number of integrals in every summand corresponds to a unique subset $e \subset [d]$. The sign in front is given by $(-1)^{|e|}$.

The following theorem states an upper bound for the error of best approximation of multivariate mixed Sobolev functions with respect to anisotropic polynomials. It turns out that $P_r f$ from (2.8) provides a good approximation of $f \in W_p^r(Q)$.

Theorem 2.7 Let $1 \leq p \leq \infty$, $r \in \mathbb{N}^d$. Then there is a constant C depending only on r, d such that for every $f \in W_p^r(Q)$

$$E_r(f)_{p, Q} \leq C \sum_{e \subset [d], e \neq \emptyset} \prod_{i \in e} \delta_i^{r_i} \|f^{(r(e))}\|_{p, Q},$$

where $\delta = \delta(Q)$ is given as in Theorem 1.1.

Proof. For simplicity we prove the theorem for the case $d = 2$ and $Q = [0, b_1] \times [0, b_2]$. Let now $f \in W_p^r(Q)$ be a bivariate function. By Hölder's and triangle inequality we obtain from (2.8) the following estimate

$$\|f - P_r f\|_{p,Q} \ll b_2^{r_2} \|f^{(0,r_2)}\|_{p,Q} + b_1^{r_1} \|f^{(r_1,0)}\|_{p,Q} + b_1^{r_1} b_2^{r_2} \|f^{(r)}\|_{p,Q}. \quad (2.16)$$

For the general case ($d > 2$) one has to take Remark 2.6 into account. \square

3 Johnen type inequalities for mixed K-functionals

For $r \in \mathbb{N}^d$, the *mixed K-functional* $K_r(f, t)_{p,Q}$ is defined for functions $f \in L_p(Q)$ and $t \in \mathbb{R}_+^d$ by

$$K_r(f, t)_{p,Q} := \inf_{g \in W_p^r(Q)} \{ \|f - g\|_{p,Q} + \sum_{e \subset [d], e \neq \emptyset} \left(\prod_{i \in e} t_i \right) \|g^{(r(e))}\|_{p,Q} \}.$$

The following technical lemma needs a further notation. Let us assume $a_i \leq c_i < d_i \leq b_i$ for $i \in [d]$. We put $I^i = [a_i, b_i]$, $I_1^i = [a_i, d_i]$, and $I_0^i = [c_i, b_i]$ and further

$$Q_e := \prod_{i=1}^d I_{\chi_e(i)}^i, \quad (3.1)$$

where χ_e denotes the characteristic function of the set $e \subset [d]$.

Lemma 3.1 *Let $1 \leq p \leq \infty$ and $r \in \mathbb{N}^d$. Then for any $f \in L_p(Q)$ the inequality*

$$K_r(f, t^r)_{p,Q} \leq C \sum_{e \subset [d]} K_r(f, t^r)_{p,Q_e}$$

holds true for all $t \in \mathbb{R}_+^d$ with $t_i \leq d_i - c_i$, $i \in [d]$. The constant C only depends on r and d .

Proof. The proof is based on an iterative argument. The first step is to observe

$$\begin{aligned} Q &= Q_1 \cup Q_0 \\ &= \left(I_1^1 \times \prod_{i \in [d] \setminus \{1\}} I^i \right) \cup \left(I_0^1 \times \prod_{i \in [d] \setminus \{1\}} I^i \right) \end{aligned}$$

and to show that

$$K_r(f, t^r)_{p,Q} \ll K_r(f, t^r)_{p,Q_1} + K_r(f, t^r)_{p,Q_0}. \quad (3.2)$$

We start with an increasing function $\varphi \in C^\infty(\mathbb{R})$ such that

$$\varphi(s) = \begin{cases} 0 & : s < 0 \\ 1 & : s > 1 \end{cases}.$$

Putting $h = d_1 - c_1$ and

$$\lambda(s) = \varphi\left(\frac{s - c_1}{h}\right), \quad s \in \mathbb{R},$$

we obtain a $C^\infty(\mathbb{R})$ -function λ that equals zero on $[a_1, c_1]$, equals one on $[d_1, b_1]$, and is increasing on $[c_1, d_1]$. As a direct consequence we get

$$\|\lambda^{(k)}\|_{\infty, \mathbb{R}} \leq h^{-k} \|\varphi^{(k)}\|_{\infty, \mathbb{R}}, \quad k \in \mathbb{N}.$$

Let now $f \in W_p^r(Q)$ and $t \in \mathbb{R}_+^d$ with $t_i \leq d_i - c_i$, $i \in [d]$. For arbitrary $g_1 \in W_p^r(Q_1)$ and $g_0 \in W_p^r(Q_0)$, put

$$\begin{aligned} g(x) &= \lambda(x_1)g_0(x) + (1 - \lambda(x_1))g_1(x) \\ &= g_1(x) + \lambda(x_1)(g_0(x) - g_1(x)). \end{aligned}$$

First of all, the function g is defined on $Q_0 \cap Q_1 \subset Q$. We extend g by g_0 on $Q_0 \setminus Q_1$ and by g_1 on $Q_1 \setminus Q_0$ and denote the result also by g . By the construction of λ this g belongs to $W_p^r(Q)$ and we have

$$\begin{aligned} \|f - g\|_{p, Q} &\leq \|\lambda(x_1)f(x) - \lambda(x_1)g_0(x) + (1 - \lambda(x_1))f(x) - (1 - \lambda(x_1))g_1(x)\|_{p, Q} \\ &\leq \|f - g_0\|_{p, Q_0} + \|f - g_1\|_{p, Q_1}. \end{aligned} \quad (3.3)$$

Furthermore, for any non-empty fixed subset $e \subset [d]$ we have

$$g^{(r(e))}(x) = g_1^{(r(e))}(x) + \sum_{k=0}^{r_1} \binom{r_1}{k} \lambda^{(r_1-k)}(x_1) (g_1^{(k, \tilde{r}(e))}(x) - g_0^{(k, \tilde{r}(e))}(x))$$

on $Q_0 \cap Q_1$, where $\tilde{r}(e)$ denotes the vector $r(e \setminus \{1\})$. Hence, for any non-empty fixed subset $e \subset [d]$ we obtain

$$\begin{aligned} &\left(\prod_{i \in e} t_i^{r_i} \right) \|g^{(r(e))}\|_{p, Q_0 \cap Q_1} \\ &\ll \left(\prod_{i \in e} t_i^{r_i} \right) \left(\|g_1^{(r(e))}\|_{p, Q_0 \cap Q_1} + \max_{0 \leq k \leq r_1} h^{-(r_1-k)} \|g_1^{(k, \tilde{r}(e))} - g_0^{(k, \tilde{r}(e))}\|_{p, Q_1 \cap Q_0} \right) \\ &\ll \left(\prod_{i \in e \setminus \{1\}} t_i^{r_i} \right) \left(t_1^{r_1 \chi_e(1)} \|g_1^{(r(e))}\|_{p, Q_0 \cap Q_1} + \max_{0 \leq k \leq r_1} \left(\frac{t_1}{h} \right)^{r_1-k} t_1^k \|g_1^{(k, \tilde{r}(e))} - g_0^{(k, \tilde{r}(e))}\|_{p, Q_1 \cap Q_0} \right). \end{aligned} \quad (3.4)$$

We apply Lemma 2.1 together with Lemma 2.2 to obtain

$$t_1^k \|g_1^{(k, \tilde{r}(e))} - g_0^{(k, \tilde{r}(e))}\|_{p, Q_0 \cap Q_1} \ll \|g_1^{(0, \tilde{r}(e))} - g_0^{(0, \tilde{r}(e))}\|_{p, Q_0 \cap Q_1} + t_1^{r_1} \|g_1^{(r(e))} - g_0^{(r(e))}\|_{p, Q_0 \cap Q_1}.$$

Plugging this into (3.4) and taking $t_1 \leq h$ into account gives in case $\tilde{r}(e) \neq 0$

$$\begin{aligned} \left(\prod_{i \in e} t_i^{r_i} \right) \|g^{(r(e))}\|_{p, Q_0 \cap Q_1} &\ll \left(\prod_{i \in e} t_i^{r_i} \right) \|g_0^{(r(e))}\|_{p, Q_0} + \left(\prod_{i \in e \setminus \{1\}} t_i^{r_i} \right) \|g_0^{(0, \tilde{r}(e))}\|_{p, Q_0} \\ &\quad + \left(\prod_{i \in e} t_i^{r_i} \right) \|g_1^{(r(e))}\|_{p, Q_1} + \left(\prod_{i \in e \setminus \{1\}} t_i^{r_i} \right) \|g_1^{(0, \tilde{r}(e))}\|_{p, Q_1}, \end{aligned} \quad (3.5)$$

and in case $\tilde{r}(e) = 0$, i.e., $e = \{1\}$,

$$\begin{aligned} \left(\prod_{i \in e} t_i^{r_i} \right) \|g^{(r(e))}\|_{p, Q_0 \cap Q_1} &\ll \left(\prod_{i \in e} t_i^{r_i} \right) \|g_0^{(r(e))}\|_{p, Q_0} + \|f - g_0\|_{p, Q_0} \\ &\quad + \left(\prod_{i \in e} t_i^{r_i} \right) \|g_1^{(r(e))}\|_{p, Q_1} + \|f - g_1\|_{p, Q_1}. \end{aligned} \quad (3.6)$$

Using that

$$\|g^{(r(e))}\|_{p,Q} \leq \|g^{(r(e))}\|_{p,Q_0 \cap Q_1} + \|g_0^{(r(e))}\|_{p,Q_0} + \|g_1^{(r(e))}\|_{p,Q_1},$$

we obtain together with (3.3), (3.5), and (3.6) the relation

$$K_r(f, t^r)_{p,Q} \ll K_r(f, t^r)_{p,Q_0} + K_r(f, t^r)_{p,Q_1}$$

which is (3.2). We continue with the same procedure, this time with Q_1 and Q_0 instead of Q , proving that (analogously for Q_1)

$$K_r(f, t^r)_{p,Q_0} \ll K_r(f, t^r)_{p,Q_{01}} + K_r(f, t^r)_{p,Q_{00}},$$

where

$$Q_{00} = \left(I_0^1 \times I_0^2 \times \prod_{i \in [d] \setminus \{1,2\}} I^i \right) \quad \text{and} \quad Q_{01} = \left(I_0^1 \times I_1^2 \times \prod_{i \in [d] \setminus \{1,2\}} I^i \right),$$

and so forth. An iteration of this argument finishes the proof. \square

3.1 Proof of Theorem 1.2

Proof. The first inequality in (1.3) follows from the definition. Namely, if $f \in L_p(Q)$, for any non-empty $e \subset [d]$ and any $g \in W_p^r(Q)$, we have

$$\begin{aligned} \omega_{r(e)}(f, t)_{p,Q} &\leq \omega_{r(e)}(f - g, t)_{p,Q} + \omega_{r(e)}(g, t)_{p,Q} \\ &\leq \left(\prod_{i \in e} 2^{r_i} \right) \left\{ \|f - g\|_{p,Q} + \left(\prod_{i \in e} t_i^{r_i} \right) \|g^{(r(e))}\|_{p,Q} \right\}. \end{aligned}$$

Indeed, the last inequality follows from the well-known relation

$$\|\Delta_h^m g\|_{p,I} \leq 2^m |h|^m \|g^{(m)}\|_{p,I}$$

for univariate functions $g \in W_p^m(I)$, which is a simple consequence of the univariate Taylor formula (2.15) and the fact that $\Delta_h^m p \equiv 0$ for a univariate polynomial of degree less than m . We iterate this relation for any index in $i \in e$ using that for frozen variables $x_1, \dots, x_{i-1}, x_{i+1}, \dots, x_d$ the univariate trace function $f(x_1, \dots, x_{i-1}, \cdot, x_{i+1}, \dots, x_d)$ belongs to the Sobolev space $W_p^{r_i}(I_i)$, see Lemma 2.2. This proves the first inequality in (1.3). Let us prove the second one. For simplicity we prove it for $d = 2$ and $t \in \mathbb{R}_+^2$, $t > 0$. If k is a natural number, then for univariate functions φ on the interval $[a, b]$, we define the operator $P_t^k, t \geq 0$, by

$$P_t^k(\varphi, x) := \varphi(x) + (-1)^{k+1} \int_{-\infty}^{\infty} \Delta_{th}^k(\varphi, x) M_k(h) dh,$$

where M_k is the B-spline of order k with knots at the integer points $0, \dots, k$, and support $[0, k]$. The function $P_t^k(\varphi)$ is defined on $[a, b - h/4]$ for $t \leq \bar{t} := h/4k^2$, where $h := b - a$. We have, see [8, page 177],

$$\{P_t^k(\varphi)\}^{(k)}(x) = t^{-k} \sum_{j=1}^k (-1)^{j+1} j^{-k} \Delta_{jt}^k(\varphi, x). \quad (3.7)$$

Put $h_i := b_i - a_i$ and $c_i := a_i + h_i/4$, $d_i := b_i - h_i/4$, $i \in [2]$. It holds $a_i < c_i < d_i < b_i$, and we will use the notation Q_e given in (3.1) for any $e \subset [d]$. In particular, we have $Q_{[2]} = [a_1, d_1] \times [a_2, d_2]$. For functions f on the parallelepiped $Q = [a_1, b_1] \times [a_2, b_2]$ the operator P_t^r , $t \in \mathbb{R}_+^2$, is defined by

$$P_t^r(f) := \prod_{i=1}^2 P_{t_i, i}^{r_i}(f),$$

where the univariate operator $P_{t_i, i}^{r_i}$ is applied to the univariate function f by considering f as a function of variable x_i with the remaining variables fixed. The function $P_t^k(f)$ is defined on $Q_{[2]}$ for $t \leq \bar{t}$, where $\bar{t}_i := h_i/4r_i^2$. We have

$$\begin{aligned} P_t^r(f, x) &= f(x) + (-1)^{r_1+1} \int_{-\infty}^{\infty} \Delta_{th}^{r_1}(f, x) M_{r_1}(h_1) dh_1 + (-1)^{r_2+1} \int_{-\infty}^{\infty} \Delta_{th}^{r_2}(f, x) M_{r_2}(h_2) dh_2 \\ &\quad + (-1)^{r_1+r_2+2} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \Delta_{th}^r(f, x) M_{r_1}(h_1) M_{r_2}(h_2) dh_1 dh_2, \end{aligned}$$

where $r^1 := (r_1, 0)$ and $r^2 := (0, r_2)$. Let us define the function $g_t = P_t^r(f)$. If $f \in L_p(Q)$, by Minkowski's inequality and properties of the B-spline M_{r_i} we get

$$\|f - g_t\|_{p, Q_{[2]}} \ll \omega_{r^1}(f, t)_{p, Q_{[2]}} + \omega_{r^2}(f, t)_{p, Q_{[2]}} + \omega_r(f, t)_{p, Q_{[2]}} = \Omega_r(f, t)_{p, Q_{[2]}}. \quad (3.8)$$

Further, by (3.7) we obtain

$$g_t^{(r^1)} = P_{t_2, 2}^{r_2}(\{P_{t_1, 1}^{r_1}(f)\}^{(r^1)}) = P_{t_2, 2}^{r_2} \left\{ t_1^{-r_1} \sum_{j_1=1}^{r_1} (-1)^{j_1+1} j_1^{-r_1} \binom{r_1}{j_1} \Delta_{j_1 t_1, 1}^{r_1}(f) \right\}.$$

Since $P_{t_2, 2}^{r_2}$ is a linear bounded operator from $L_p(Q_{[2]})$ into $L_p(Q_{[2]})$ and further $\|\Delta_{j_1 t_1, 1}^{r_1}(f)\|_{p, Q_{[2]}} \ll \omega_{r^1}(f, t)_{p, Q_{[2]}}$, we have

$$t_1^{r_1} \|g_t^{(r^1)}\|_{p, Q_{[2]}} \ll \omega_{r^1}(f, t)_{p, Q_{[2]}}. \quad (3.9)$$

Similarly, we can prove that

$$t_2^{r_2} \|g_t^{(r^2)}\|_{p, Q_{[2]}} \ll \omega_{r^2}(f, t)_{p, Q_{[2]}}.$$

Again, by (3.7) we get

$$g_t^{(r)} = t_1^{-r_1} t_2^{-r_2} \sum_{j_1=1}^{r_1} \sum_{j_2=1}^{r_2} (-1)^{j_1+j_2+2} j_1^{-r_1} j_2^{-r_2} \binom{r_1}{j_1} \binom{r_2}{j_2} \Delta_{j_1 j_2}^r(f).$$

From the inequality $\|\Delta_{j_1 j_2}^r(f)\|_{p, Q} \ll \omega_r(f, t)_{p, Q_{[2]}}$ it follows that

$$t_1^{r_1} t_2^{r_2} \|g_t^{(r)}\|_{p, Q_{[2]}} \ll \omega_r(f, t)_{p, Q_{[2]}}. \quad (3.10)$$

Combining (3.8), (3.9)–(3.10) gives

$$\|f - g_t\|_{p, Q_{[2]}} + t_1^{r_1} \|g_t^{(r^1)}\|_{p, Q_{[2]}} + t_2^{r_2} \|g_t^{(r^2)}\|_{p, Q_{[2]}} + t_1^{r_1} t_2^{r_2} \|g_t^{(r)}\|_{p, Q_{[2]}} \ll \Omega_r(f, t)_{p, Q}.$$

Therefore, we get

$$K_r(f, t^r)_{p, Q_{[2]}} \ll \Omega_r(f, t)_{p, Q},$$

and in a similar way

$$K_r(f, t^r)_{p, Q_e} \ll \Omega_r(f, t)_{p, Q}$$

for any subset $e \subset [2]$, where Q_e is given by (3.1). The last inequality and Lemma 3.1 prove (1.3) for $t \leq \bar{t}$. Now take a function $\bar{g} \in W_p^r(Q)$ such that

$$\|f - \bar{g}\|_{p, Q} + \bar{t}_1^{r_1} \|\bar{g}^{(r_1)}\|_{p, Q} + \bar{t}_2^{r_2} \|\bar{g}^{(r_2)}\|_{p, Q} + \bar{t}_1^{r_1} \bar{t}_2^{r_2} \|\bar{g}^{(r)}\|_{p, Q} \ll \Omega_r(f, \bar{t})_{p, Q}. \quad (3.11)$$

By Theorem 2.7 we have

$$\|\bar{g} - T_r(\bar{g})\|_{p, Q} \ll \bar{t}_1^{r_1} \|\bar{g}^{(r_1)}\|_{p, Q} + \bar{t}_2^{r_2} \|\bar{g}^{(r_2)}\|_{p, Q} + \bar{t}_1^{r_1} \bar{t}_2^{r_2} \|\bar{g}^{(r)}\|_{p, Q}. \quad (3.12)$$

Since $T_r(\bar{g}) \in W_p^r(Q)$ and $(T_r(\bar{g}))^{(r(e))} = 0$ for every non-empty subset $e \subset [d]$, it holds for all $t > \bar{t}$

$$\begin{aligned} K_r(f, t^r)_{p, Q} &\leq \|f - T_r(\bar{g})\|_{p, Q} \\ &\leq \|f - \bar{g}\|_{p, Q} + \|\bar{g} - T_r(\bar{g})\|_{p, Q} \\ &\ll \Omega_r(f, \bar{t})_{p, Q} \leq \Omega_r(f, t)_{p, Q}, \end{aligned}$$

where the third step combines (3.11) and (3.12). Therefore, (1.3) has been proved for arbitrary $t > 0$. \square

4 Whitney type inequalities

Using the results from Section 3 we are now able to prove Theorem 1.1.

Proof. The first inequality in (1.2) is trivial. Indeed, if $f \in L_p(Q)$ then for any non-empty $e \subset [d]$ and any $\varphi \in \mathcal{P}_r$ we have

$$\begin{aligned} \omega_{r(e)}(f, \delta)_{p, Q} &= \omega_{r(e)}(f - \varphi, \delta)_{p, Q} \\ &\leq \left(\prod_{i \in e} 2^{r_i} \right) \|f - \varphi\|_{p, Q}. \end{aligned}$$

Hence, we obtain the first inequality in (1.2). On the other hand, from Theorem 2.7 it follows that for any $g \in W_p^r(Q)$

$$\begin{aligned} E_r(f)_{p, Q} &\leq \|f - g\|_{p, Q} + E_r(g)_{p, Q} \\ &\leq \|f - g\|_{p, Q} + \|g - T_r(g)\|_{p, Q} \\ &\ll \|f - g\|_{p, Q} + \left(\prod_{i \in e} \delta_i^{r_i} \right) \|g^{(r(e))}\|_{p, Q}. \end{aligned}$$

Hence, we get

$$E_r(f)_{p, Q} \ll K_r(f, \delta^r)_{p, Q}.$$

By Theorem 1.2 we have proved the second inequality in (1.2). \square

The result in Theorem 1.1 can be slightly modified. For $r \in \mathbb{Z}_+^d$, $h \in \mathbb{R}^d$, $e \subset [d]$ and a d -variate function $f : \mathbb{R}^d \rightarrow \mathbb{R}$ the mixed p -mean modulus of smoothness of order $r(e)$ is given by

$$w_{r(e)}(f, t)_p := \left(\left(\prod_{i \in e} t_i^{-1} \right) \int_{U(t)} \int_{Q_{r(e)h}} |\Delta_h^{r(e)}(f, x)|^p dx dh \right)^{1/p}, \quad t \in \mathbb{R}_+^d,$$

where $U(t) := \{h \in \mathbb{R}^d : |h_i| \leq t_i, i \in [d]\}$, with the usual change of the outer mean integral to sup if $p = \infty$. This leads to the definition of the *total mixed p -mean modulus of smoothness of order r* for $r \in \mathbb{N}^d$ by

$$W_r(f, t)_{p,Q} := \sum_{e \subset [d], e \neq \emptyset} w_{r(e)}(f, t)_{p,Q}, \quad t \in \mathbb{R}_+^d.$$

Note that $W_r(f, t)_{p,Q}$ coincides with $\Omega_r(f, t)_{p,Q}$ when $p = \infty$. In a way similar to the proof of Theorem 1.1 we can prove the following result.

Theorem 4.1 *Let $1 \leq p \leq \infty$, $r \in \mathbb{N}^d$. Then there are constants C, C' depending only on r, d such that for every $f \in L_p(Q)$*

$$CW_r(f, \delta)_{p,Q} \leq E_r(f)_{p,Q} \leq C'W_r(f, \delta)_{p,Q}$$

where $\delta = \delta(Q)$.

Remark 4.2 *A corresponding inequality in the case $0 < p < 1$ is so far left open for subsequent contributions. It seems that the modulus $W_r(f, t)_{p,Q}$ is suitable to treat this case, cf. the appendix of [12].*

Acknowledgements Since main parts of the paper have been worked out during a stay of the second named author at the Information Technology Institute of the Vietnam National University, Hanoi in 2010, Tino Ullrich would like to thank here Prof. Dinh Dung for his kind invitation and warm hospitality. He would also like to thank the Hausdorff-Center for Mathematics at University of Bonn (Germany) and Prof. Holger Rauhut for giving additional financial support. The work of the first named author was supported by the National Foundation for Development of Science and Technology (Vietnam). Both authors would also like to thank Dany Leviatan and two unknown referees for a critical reading of the manuscript and for several valuable comments how to improve it.

References

- [1] O.V. Besov, V.P. Il'in, and S.M. Nikol'skij, Integral representations of functions and embedding theorems, Vol. 1., V.H. Winston & Sons, Washington D.C., 1978.
- [2] Yu.A. Brudnyĭ, On a theorem on best local approximations, *Kazanskii Gosudarstvennyi Universitet. Uchenye Zapiski* **124**, No. 6 (1964), 43–49.

- [3] Yu.A. Brudnyi, A multidimensional analogue of a certain theorem of Whitney, *Math. USSR-Sb.* **2** (1970), 157–170.
- [4] Yu.A. Brudnyi, Approximation of functions of n variables by quasipolynomials, *Izv. Akad. Nauk SSSR Ser. Mat.* **34**, No. 3 (1970), 564–583.
- [5] O.V. Davydov, Sequences of rectangular Fourier sums of continuous functions with given majorants of the mixed moduli of smoothness, *Mat. Sb.* **187**(1996), No7, 35–58.
- [6] S. Dekel, On the equivalence of the modulus of smoothness and the K-functional over convex domains, *Journ. Approx. Theory* **162** (2010), 349–362.
- [7] S. Dekel and D. Leviatan, Whitney estimates for convex domains with applications to multivariate piecewise polynomial approximation, *Found. Comput. Math.* **4** (2004), 345–368.
- [8] R.A. DeVore and G.G. Lorentz, *Constructive approximation*, Springer-Verlag, New York, 1993.
- [9] R.A. DeVore, P.P. Petrushev, and V.N. Temlyakov, Multivariate trigonometric polynomial approximations with frequencies from the hyperbolic cross, *Mat. Zametki* **56**, No. 3 (1994), 36–63.
- [10] Dinh Dung, Approximation of functions of several variables on tori by trigonometric polynomials, *Mat. Sb.* **131**(1986), 251-271.
- [11] Dinh Dung, B-spline quasi-interpolant representations and sampling recovery of functions with mixed smoothness, *Journal of Complexity* (2011), doi:10.1016/j.jco.2011.02.004.
- [12] L. I. Hedberg and Yu. Netrusov, An axiomatic approach to function spaces, spectral synthesis, and Luzin approximation, *Mem. Amer. Math. Soc.* **188**, No. 882 (2007).
- [13] G. Garrigós and A. Tabacco, Wavelet decompositions of anisotropic Besov spaces, *Math. Nachr.* **239–240** (2002), 80–102.
- [14] H. Johnen, Inequalities connected with moduli of smoothness, *Math. Vesnik* **3** (1972), 289–303.
- [15] H. Johnen and K. Scherer, On the equivalence of the K-functional and moduli of continuity and some applications, in: *Constructive Theory of Functions of Several Variables*, Proc. Conf. Math. Res. Inst., Oberwolfach, 1976, 119–140; Lecture Notes **571**, Springer Berlin, 1977.
- [16] J. Peetre, A theory of interpolation of normed spaces, Notes Universidade de Brasilia, 1963.
- [17] G. Sparr, Interpolation of several Banach spaces, *Ann. Mat. Pur. Appl.* **99**, No. 1 (1974), 247-316.
- [18] È.A. Storozhenko, Approximation by algebraic polynomials of functions in the class L^p , $0 < p < 1$, *Izv. Akad. Nauk SSSR, Ser. Mat.* **41** (1977), 652–662.
- [19] È.A. Storozhenko and P. Oswald, Jackson’s theorem in the spaces $L^p(\mathbb{R}^k)$, $0 < p < 1$, *Siberian Math. J.* **19** (1978), 630-640.

- [20] H.-J. Schmeisser and H. Triebel, Topics in Fourier analysis and function spaces, Wiley, Chichester, 1987.
- [21] H. Triebel, Bases in function spaces, sampling, discrepancy, numerical integration, EMS Tracts in Mathematics, Vol. 11, European Mathematical Society (EMS), Zürich, 2010.
- [22] T. Ullrich, Function spaces with dominating mixed smoothness, characterization by differences, *Jenaer Schriften zur Mathematik und Informatik*, Math/Inf/05/06 (2006).
- [23] J. Vybiral, Function spaces with dominating mixed smoothness, *Dissertationes Math.* **436** (2006), 73 pp.
- [24] H. Whitney, On functions with bounded n th difference, *J. Math. Pures Appl.* **36** (1957), 67–95.